
antibodies -online.com







anti-RAP2A antibody (AA 124-159)



Image



\sim	
()\/\	rview
\cup	1 410 44

Quantity:	200 μL
Target:	RAP2A
Binding Specificity:	AA 124-159
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAP2A antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This RAP2A antibody is generated from a rabbit immunized with a KLH conjugated synthetic

Immunogen:	This RAP2A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 124-159 amino acids from human RAP2A.
Clone:	RB53398
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	RAP2A
Alternative Name:	RAP2A (RAP2A Products)
Background:	Small GTP-binding protein which cycles between a GDP- bound inactive and a GTP-bound

active form. In its active form interacts with and regulates several effectors including MAP4K4, MINK1 and TNIK. Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development. More generally, it is part of several signaling cascades and may regulate cytoskeletal rearrangements, cell migration, cell adhesion and cell spreading.

Molecular Weight:

20615

UniProt:

P10114

Application Details


Application Notes: WB: 1:2000

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Images



Western Blotting

Image 1. Anti-RA Antibody (C-Term) at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.