

Datasheet for ABIN6242630
anti-SMAD1 antibody (AA 20-330)



[Go to Product page](#)

4 Images

Overview

Quantity:	200 µL
Target:	SMAD1
Binding Specificity:	AA 20-330
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SMAD1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This SMAD1 antibody is generated from a mouse immunized with a recombinant protein between 20-330 amino acids from human SMAD1.
Clone:	1356CT119-18-55
Isotype:	IgG1 kappa
Purification:	This antibody is purified through a protein G column, followed by dialysis against PBS.

Target Details

Target:	SMAD1
Alternative Name:	SMAD1 (SMAD1 Products)

Target Details


Background:	Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD1 is a receptor-regulated SMAD (R-SMAD). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. May act synergistically with SMAD4 and YY1 in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression.
Molecular Weight:	52260
UniProt:	Q15797
Pathways:	Stem Cell Maintenance , Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development

Application Details

Application Notes:	IF: 1:25. WB: 1:2000. IHC-P: 1:25. IHC-P: 1:25
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified monoclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. (ABIN6242630 and ABIN6577239) staining SD1 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hour at 37 °C. An undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunofluorescence

Image 2. Immunofluorescent analysis of 4 % paraformaldehyde-fixed, 0.1 % Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling SD1 with (ABIN6242630 and ABIN6577239) at 1/25 dilution, followed by DyLight® 488-conjugated goat anti-mouse IgG (NH174309) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing nucleus and weak cytoplasm staining on HeLa cell line. Cytoplasmic actin is detected with DyLight® 554 Phalloidin (OI17558410) at 1/100 dilution (red). The nuclear counterstain is DAPI (blue).



Western Blotting

Image 3. All lanes : Anti-SD1 Antibody at 1:2000 dilution
Lane 1: HT-1080 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 60 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6242630.