antibodies - online.com







anti-SRY antibody (AA 81-112)





O	:
1 1\/\pir	view
\circ	V I C V V

Overview	
Quantity:	400 μL
Target:	SRY
Binding Specificity:	AA 81-112
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SRY antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This SRY antibody is generated from a rabbit immunized with a KLH conjugated synthetic
	peptide between 81-112 amino acids from the Central region of human SRY.
Clone:	RB53826
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	SRY
Alternative Name:	SRY (SRY Products)
Background:	Transcriptional regulator that controls a genetic switch in male development. It is necessary

Target Details

and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (By similarity). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene regulation including promoter activation or repression (By similarity). Promotes DNA bending. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Also involved in pre-mRNA splicing. Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3'.

Molecular Weight:

23884

UniProt:

Q05066

6 months

Application Details

Application Notes: WB: 1:1000


Restrictions: For

For Research Use only

Handling

Expiry Date:

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C



Western Blotting

Image 1. Anti-SRY Antibody (Center)at 1:1000 dilution + K562 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 24 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.