antibodies -online.com





anti-UGT2B4 antibody (AA 338-370)





Go to Product page

\sim			
	N/6	1//r	$I \cap V$

Overview		
Quantity:	200 μL	
Target:	UGT2B4	
Binding Specificity:	AA 338-370	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UGT2B4 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This UGT2B4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic	
	peptide between 338-370 amino acids from the Central region of human UGT2B4.	
Clone:	RB53182	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	UGT2B4	
Alternative Name:	UGT2B4 (UGT2B4 Products)	
Background:	UDPGTs are of major importance in the conjugation and subsequent elimination of potentially	

Target Details

Molecular Weight: 60513

UniProt: P06133

Pathways: Steroid Hormone Biosynthesis


Application Details

Application Notes: WB: 1:2000. WB: 1:2000. WB: 1:2000

Restrictions: For Research Use only


Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	




Western Blotting

Image 1. Anti-UGT2B4 Antibody (Center)at 1:2000 dilution + human liver lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 2. Anti-UGT2B4 Antibody (Center)at 1:2000 dilution + human cerebellum lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. Anti-UGT2B4 Antibody (Center)at 1:2000 dilution + human liver lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Please check the product details page for more images. Overall 4 images are available for ABIN6242687.