antibodies -online.com







anti-NCOA2 antibody (AA 706-739)



Image



\sim	
()\/\Di	view
	VICVV

Quantity:	200 μL
Target:	NCOA2
Binding Specificity:	AA 706-739
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NCOA2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This NCOA2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic
	peptide between 706-739 amino acids from the Central region of human NCOA2.
Clone:	RB54073
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	NCOA2
Alternative Name:	NCOA2 (NCOA2 Products)

steroid binding domain (AF- 2) but not of the modulating N-terminal domain (AF-1). Required with NCOA1 to control energy balance between white and brown adipose tissues. Critical regulator of glucose metabolism regulation, acts as RORA coactivator to specifically modulate G6PC expression. Involved in the positive regulation of the transcriptional activity of the glucocorticoid receptor NR3C1 by sumoylation enhancer RWDD3. Positively regulates the circadian clock by acting as a transcriptional coactivator for the CLOCK- ARNTL/BMAL1 heterodimer (By similarity).

Molecular Weight:

159157

UniProt:

015596

Pathways:

Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Nuclear Hormone Receptor Binding, Regulation of Lipid Metabolism by PPARalpha, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling


Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months



Western Blotting

Image 1. Anti-NCOA2 Antibody (Center) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 159 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.