

Datasheet for ABIN6242969
anti-NMNAT1 antibody (AA 200-232)[Go to Product page](#)

2 Images

Overview

Quantity:	200 µL
Target:	NMNAT1
Binding Specificity:	AA 200-232
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NMNAT1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This NMNAT1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 200-232 amino acids from human NMNAT1.
Clone:	RB55363
Isotype:	Ig Fraction
Predicted Reactivity:	B, M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	NMNAT1
Alternative Name:	NMNAT1 (NMNAT1 Products)

Target Details

Background: Catalyzes the formation of NAD(+) from nicotinamide mononucleotide (NMN) and ATP. Can also use the deamidated form, nicotinic acid mononucleotide (NaMN) as substrate with the same efficiency. Can use triazofurin monophosphate (TrMP) as substrate. Also catalyzes the reverse reaction, i.e. the pyrophosphorolytic cleavage of NAD(+). For the pyrophosphorolytic activity, prefers NAD(+) and NAAD as substrates and degrades NADH, nicotinic acid adenine dinucleotide phosphate (NHD) and nicotinamide guanine dinucleotide (NGD) less effectively. Fails to cleave phosphorylated dinucleotides NADP(+), NADPH and NAADP(+). Protects against axonal degeneration following mechanical or toxic insults.

Molecular Weight: 31932

UniProt: [Q9HAN9](#)

Application Details

Application Notes: WB: 1:2000. WB: 1:2000

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-20 °C

Expiry Date: 6 months



Western Blotting

Image 1. Anti-NMNAT1 Antibody (C-Term) at 1:2000 dilution + human heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 2. Anti-NMNAT1 Antibody (C-Term) at 1:2000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.