

Datasheet for ABIN6243115
anti-CKS2 antibody (AA 6-40)[Go to Product page](#)

2 Images

Overview

Quantity:	200 µL
Target:	CKS2
Binding Specificity:	AA 6-40
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CKS2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This CKS2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 6-40 amino acids from the human region of human CKS2.
Clone:	RB30893
Isotype:	Ig Fraction
Predicted Reactivity:	B, M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CKS2
Alternative Name:	CKS2 (CKS2 Products)

Target Details

Background: Binds to the catalytic subunit of the cyclin dependent kinases and is essential for their biological function.

Molecular Weight: 9860

UniProt: [P33552](#)

Application Details

Application Notes: WB: 1:2000. FC: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Expiry Date: 6 months

Images



Flow Cytometry

Image 1. Overlay histogram showing HeLa cells stained with (ABIN6243115 and ABIN6578863)(green line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN6243115 and ABIN6578863), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was rabbit IgG1 (1 µ



g/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Western Blotting

Image 2. All lanes : Anti-CKS2 Antibody (N-Term) at 1:2000 dilution Lane 1: Caco2 whole cell lysate Lane 2: RI 8226 whole cell lysate Lane 3: SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 10 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.