antibodies -online.com





anti-POLR2H antibody (N-Term)

2 Images



Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Overview		
Quantity:	400 μL	
Target:	POLR2H	
Binding Specificity:	AA 16-50, N-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This POLR2H antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This POLR2H antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 16-50 amino acids from the N-terminal region of human POLR2H.	
Clone:	RB49268	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	POLR2H	
Alternative Name:	POLR2H (POLR2H Products)	
Background:	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four	

ribonucleoside triphosphates as substrates. Common component of RNA polymerases I, II and III which synthesize ribosomal RNA precursors, mRNA precursors and many functional non-coding RNAs, and small RNAs, such as 5S rRNA and tRNAs, respectively.

Molecular Weight: 17143
UniProt: P52434

Pathways: Regulatory RNA Pathways

Application Details


Application Notes: WB: 1:1000. WB: 1:1000

Restrictions: For Research Use only

Handling


Format:	Liquid		
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	4 °C,-20 °C		
Expiry Date:	6 months		

Images



Western Blotting

Image 1. Western blot analysis of lysate from mouse spleen tissue lysate, using POLR2H Antibody (N-term) (ABIN6243172 and ABIN6577586). (ABIN6243172 and ABIN6577586) was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35 μ g.



Western Blotting

Image 2. Western blot analysis of lysate from A431 cell line, using POLR2H Antibody (N-term) (ABIN6243172 and ABIN6577586). (ABIN6243172 and ABIN6577586) was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at $20 \, \mu g$.