antibodies -online.com





anti-HVEM antibody (C-Term)





Overview

Overview	
Quantity:	400 μL
Target:	HVEM (TNFRSF14)
Binding Specificity:	AA 269-302, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HVEM antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This TNFRSF14 antibody is generated from a rabbit immunized with a KLH conjugated
	synthetic peptide between 269-302 amino acids from the C-terminal region of human
	TNFRSF14.
Clone:	RB52847
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	HVEM (TNFRSF14)
Alternative Name:	TNFRSF14 (TNFRSF14 Products)

Target Details


Storage:

Expiry Date:

4 °C,-20 °C

6 months

Target Type:	Viral Protein	
Background:	Receptor for BTLA. Receptor for TNFSF14/LIGHT and homotrimeric TNFSF1/lymphotoxin-	
	alpha. Involved in lymphocyte activation. Plays an important role in HSV pathogenesis because	
	it enhanced the entry of several wild-type HSV strains of both serotypes into CHO cells, and	
	mediated HSV entry into activated human T-cells.	
Molecular Weight:	30392	
UniProt:	Q92956	
Pathways:	Production of Molecular Mediator of Immune Response, Cancer Immune Checkpoints	
Application Details		
Application Notes:	WB: 1:1000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	



Western Blotting

Image 1. Anti-TNFRSF14 Antibody (C-term)at 1:1000 dilution + Raji whole cell lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size: 30 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.