antibodies - online.com







anti-NOVA1 antibody (AA 139-173)

Images



Overview

400 μL
NOVA1
AA 139-173
Human
Rabbit
Polyclonal
This NOVA1 antibody is un-conjugated
Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
This NOVA1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic
peptide between 139-173 amino acids from the Central region of human NOVA1.
RB50876
Ig Fraction
М
This antibody is purified through a protein A column, followed by peptide affinity purification.
NOVA1

Target Details

Background:	May regulate RNA splicing or metabolism in a specific subset of developing neurons.
Molecular Weight:	51727
UniProt:	P51513


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:25
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffinembedded H. astroglioma section using NOVA1 Antibody (Center) (ABIN6243271 and ABIN6577731). (ABIN6243271 and ABIN6577731) was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Western Blotting

Image 2. Western blot analysis of lysates from A431, Hela cell line (from left to right), using NOVA1 Antibody (Center) (ABIN6243271 and ABIN6577731). (ABIN6243271 and ABIN6577731) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.