# antibodies -online.com





# anti-UHRF1 antibody (C-Term)

2 Images



Go to Product page

$\sim$						
	1//	Д	r۱	/1	$\triangle$	٨

Quantity:	400 μL	
Target:	UHRF1	
Binding Specificity:	AA 596-628, C-Term	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UHRF1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This Mouse Uhrf1 antibody is generated from a rabbit immunized with a KLH conjugated	
	synthetic peptide between 596-628 amino acids from the C-terminal region of Mouse Uhrf1.	
Clone:	RB51453	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	UHRF1	
Alternative Name:	Uhrf1 (UHRF1 Products)	
Background:	Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and	

chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD- type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. May be involved in DNA repair.

Molecular Weight:

88304

UniProt:

Q8VDF2

### **Application Details**

Application Notes:


WB: 1:1000. WB: 1:2000


Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	





#### **Western Blotting**

**Image 1.** Anti-Uhrf1 Antibody (C-term) at 1:2000 dilution + mouse thymus lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 88 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

#### **Western Blotting**

**Image 2.** Anti-Uhrf1 Antibody (C-term) at 1:1000 dilution + mouse heart lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size: 88 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.