
antibodies - online.com







anti-CNDP1 antibody (AA 128-142)

Images



\sim						
	1//	Д	r۱	/1	\triangle	٨

0.00.000		
Quantity:	400 μL	
Target:	CNDP1	
Binding Specificity:	AA 128-142	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CNDP1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This CNDP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 128-142 amino acids from the Central region of human CNDP1.	
Clone:	RB47679	
Isotype:	Ig Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	CNDP1	
Alternative Name:	CNDP1 (CNDP1 Products)	
Molecular Weight:	56706	

Target Details

UniProt:

Q96KN2

Application Details


Application Notes:	WB: 1:1000. WB: 1:1000. WB: 1:1000
--------------------	------------------------------------

Restrictions: For Research Use only

Handling


Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	


Images



Western Blotting

Image 1. Western blot analysis of lysate from human plasma tissue lysate, using CNDP1 Antibody (Center) (ABIN6243352 and ABIN6577403). (ABIN6243352 and ABIN6577403) was diluted at 1:1000. A goat anti-rabbit lgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20 μ g.





Western Blotting

Image 2. Western blot analysis of lysates from mouse kidney and rat kidney tissue lysate(from left to right), using CNDP1 Antibody (Center)(R). R was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 μ g per lane.

Western Blotting

Image 3. Western blot analysis of lysates from human plasma, mouse kidney, rat kidney tissue lysate (from left to right), using CNDP1 Antibody (Center) (ABIN6243352 and ABIN6577403). (ABIN6243352 and ABIN6577403) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 μg per lane.