

Datasheet for ABIN6243567
anti-MIF4GD antibody (AA 137-169)[Go to Product page](#)

3 Images

Overview

Quantity:	200 µL
Target:	MIF4GD
Binding Specificity:	AA 137-169
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MIF4GD antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:	This MIF4GD antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 137-169 amino acids from human MIF4GD.
Clone:	RB55240
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	MIF4GD
Alternative Name:	MIF4GD (MIF4GD Products)
Molecular Weight:	25423

Target Details

UniProt: [A9UHW6](#)

Application Details

Application Notes: IF: 1:25. WB: 1:2000. FC: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Expiry Date: 6 months

Images




Flow Cytometry

Image 1. Overlay histogram showing A431 cells stained with (ABIN6243567 and ABIN6578806)(green line). The cells were fixed with 2 % paraformaldehyde and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Western Blotting

Image 2. Anti-MIF4GD Antibody (C-Term) at 1:2000 dilution + Mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Immunofluorescence

Image 3. Immunofluorescent analysis of 4 % paraformaldehyde-fixed, 0.1 % Triton X-100 permeabilized U-2 OS (human osteosarcoma cell line) cells labeling MIF4GD with (ABIN6243567 and ABIN6578806) at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm and weak nucleus staining on U-2 OS cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DI (blue).