

Datasheet for ABIN6243590

anti-HNRNPUL2 antibody (AA 337-350)**3** Images[Go to Product page](#)

Overview

Quantity:	200 µL
Target:	HNRNPUL2
Binding Specificity:	AA 337-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPUL2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This HNRNPUL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 337-350 amino acids from the Central region of human HNRNPUL2.
Clone:	RB53002
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	HNRNPUL2
Alternative Name:	HNRNPUL2 (HNRNPUL2 Products)
UniProt:	Q1KMD3

Application Details

Application Notes: WB: 1:8000. WB: 1:2000. IHC-P: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. (ABIN6243590 and ABIN6578256) staining HNRNPUL2 in human stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hour at 37 °C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Western Blotting

Image 2. Anti-HNRNPUL2 Antibody (Center) at 1:8000 dilution + Hela whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. All lanes : Anti-HNRNPUL2 Antibody (Center) at 1:2000 dilution. Lane 1: MCF-7 whole cell lysates. Lane 2: Jurkat whole cell lysates. Lane 3: 293T/17 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.