antibodies -online.com





anti-CTR9 antibody (C-Term)





Go to Product page

\sim					
	1//	r۱.	/ I	\triangle	٨

Quantity:	400 μL		
Target:	CTR9		
Binding Specificity:	AA 1022-1056, C-Term		
Reactivity:	Human, Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This CTR9 antibody is un-conjugated		
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)		
Product Details			
Immunogen:	This Mouse Ctr9 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1022-1056 amino acids from the C-terminal region of Mouse Ctr9.		
Clone:	RB51042		
Isotype:	lg Fraction		
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.		
Target Details			
Target:	CTR9		
Alternative Name:	Ctr9 (CTR9 Products)		

Target Details

Background:

Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both indepentently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys- 4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1), UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Required for mono- and trimethylation on histone H3 'Lys- 4' (H3K4me3) and dimethylation on histone H3 'Lys-79' (H3K4me3). Required for Hox gene transcription (By similarity). Required for the trimethylation of histone H3 'Lys-4' (H3K4me3) on genes involved in stem cell pluripotency, this function is synergistic with CXXC1 indicative for an involvement of the SET1 complex. Involved in transcriptional regulation of IL6-responsive genes and in JAK-STAT pathway, may regulate DNA-association of STAT3.

Molecular Weight: 133408
UniProt: Q62018

Cellular Response to Molecule of Bacterial Origin, Stem Cell Maintenance

Application Details

Application Notes: IF: 1:25. WB: 1:1000. IHC-P: 1:25. IHC-P: 1:25

For Research Use only

Handling

Restrictions:

Pathways:

Format:

Buffer:

Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative:


Sodium azide

Precaution of Use:

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

	should be handled by trained staff only.		
Storage:	4 °C,-20 °C		
Expiry Date:	6 months		

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffinembedded M. colon section using Mouse Ctr9 Antibody (Cterm) (ABIN6243851 and ABIN6577807). (ABIN6243851 and ABIN6577807) was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunofluorescence

Image 2. Fluorescent image of A431 cells stained with Mouse Ctr9 Antibody (C-term) (ABIN6243851 and ABIN6577807). (ABIN6243851 and ABIN6577807) was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western Blotting

Image 3. Western blot analysis of lysates from Jurkat, HT-29, mouse BA/F3, mouse F9 cell line, mouse cerebellum, mouse testis tissue (from left to right), using Mouse Ctr9 Antibody (C-term) (ABIN6243851 and ABIN6577807). (ABIN6243851 and ABIN6577807) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 μg per lane.

Please check the product details page for more images. Overall 4 images are available for ABIN6243851.