antibodies - online.com







anti-TAL1 antibody (AA 154-188)

Images



\sim				
	11/6	٦r١	/10	۱۸.

Background:

Quantity:	400 μL	
Target:	TAL1	
Binding Specificity:	AA 154-188	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TAL1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This mouse Tal1 antibody is generated from a rabbit immunized with a KLH conjugated	
	synthetic peptide between 154-188 amino acids from the Central region of mouse Tal1.	
Clone:	RB50954	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	TAL1	
Alternative Name:	Tal1 (TAL1 Products)	

Implicated in the genesis of hemopoietic malignancies. It may play an important role in

Target Details

	hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation.
Molecular Weight:	34279
UniProt:	P22091
Pathways:	Stem Cell Maintenance


Application Details

Application Notes:	WB: 1:2000. WB: 1:1000
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Images



Western Blotting

Image 1. Anti-(Mouse) Tal1 Antibody (Center) at 1:1000 dilution + mouse spleen lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 34 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 2. Anti-(Mouse) Tal1 Antibody (Center) at 1:2000 dilution + K562 whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 34 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.