

Datasheet for ABIN6244086  
**anti-NAT10 antibody (AA 146-178)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	200 µL
Target:	NAT10
Binding Specificity:	AA 146-178
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NAT10 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

## Product Details

Immunogen:	This NAT10 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 146-178 amino acids from the mouse region of mouse NAT10.
Clone:	RB57638
Isotype:	Ig Fraction
Predicted Reactivity:	M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	NAT10
Alternative Name:	NAT10 ( <a href="#">NAT10 Products</a> )

## Target Details

---

**Background:** Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome de-condensation. Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization. Acetylates alpha-tubulin, which may affect microtubule stability and cell division.

**Molecular Weight:** 115730

**UniProt:** [Q9H0A0](#)

## Application Details

---

**Application Notes:** WB: 1:2000. FC: 1:25

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid


**Buffer:** Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

**Preservative:** Sodium azide

**Precaution of Use:** This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


**Storage:** 4 °C, -20 °C

**Expiry Date:** 6 months



### Flow Cytometry

**Image 1.** Overlay histogram showing HeLa cells stained with (ABIN6244086 and ABIN6578861)(green line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN6244086 and ABIN6578861), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.



### Western Blotting

**Image 2.** All lanes : Anti-NAT10 Antibody (N-Term) at 1:2000 dilution Lane 1: Human lung lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 116 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.