

Datasheet for ABIN6244171
anti-SBK1 antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

| | |
|----------------------|-------------------------------------|
| Quantity: | 400 µL |
| Target: | SBK1 |
| Binding Specificity: | AA 65-98, N-Term |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SBK1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | This Mouse Sbk1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 65-98 amino acids from the N-terminal region of Mouse Sbk1. |
| Clone: | RB44718 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | Rat |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | SBK1 |
| Alternative Name: | Sbk1 (SBK1 Products) |

Target Details

| | |
|-------------------|--|
| Background: | May be involved in signal-transduction pathways related to the control of brain development (By similarity). |
| Molecular Weight: | 45696 |
| UniProt: | Q8QZX0 |


Application Details

| | |
|--------------------|-----------------------|
| Application Notes: | WB: 1:500-1:1000 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Expiry Date: | 6 months |

Images



Western Blotting

Image 1. Western blot analysis of lysates from mouse heart, mouse brain tissue lysate (from left to right), using Sbk1 Antibody (N-term) (ABIN6244171 and ABIN6577643). (ABIN6244171 and ABIN6577643) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.