

Datasheet for ABIN6244180  
**anti-UHRF2 antibody (AA 472-505)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	80 µL
Target:	UHRF2
Binding Specificity:	AA 472-505
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UHRF2 antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	This Mouse Uhrf2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 472-505 amino acids from the Central region of Mouse Uhrf2.
Clone:	RB52404
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	UHRF2
Alternative Name:	Uhrf2 ( <a href="#">UHRF2 Products</a> )
Background:	E3 SUMO-, but not ubiquitin-, protein ligase for ZNF131 (By similarity). E3 ubiquitin-protein

## Target Details

ligase that is an intermolecular hub protein in the cell cycle network. Ubiquitinates cyclins, CCND1 and CCNE1, in an apparently phosphorylation-independent manner and induces G1 arrest. Also ubiquitinates PCNP leading to its degradation by the proteasome. Through cooperative DNA and histone binding, may contribute to a tighter epigenetic control of gene expression in differentiated cells.

Molecular Weight: 90106

UniProt: [Q7TMI3](#)

## Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

## Images



### Western Blotting

**Image 1.** Anti-Uhrf2 Antibody (Center)at 1:1000 dilution + Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 90 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.