antibodies.com

Datasheet for ABIN6244220 anti-OR2T7 antibody (C-Term)

Image



Overview

| Quantity: | 50 μL |
|----------------------|--------------------------------------|
| Target: | OR2T7 |
| Binding Specificity: | AA 264-299, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This OR2T7 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | This OR2T7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 264-299 amino acids from the C-terminal region of human OR2T7. |
|---------------|---|
| Clone: | RB47084 |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |


Target Details

| Target: | OR2T7 |
|-------------------|------------------------|
| Alternative Name: | OR2T7 (OR2T7 Products) |
| Background: | Odorant receptor. |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6244220 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

| Target Details | | |
|---------------------|--|--|
| Molecular Weight: | 34287 | |
| UniProt: | P0C7T2 | |
| Application Details | | |
| Application Notes: | WB: 1:2000 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Expiry Date: | 6 months | |

Images



Western Blotting

Image 1. All lanes : Anti-OR2T7 Antibody (C-term) at 1:2000 dilution Lane 1: human brain lysate Lane 2: HT-29 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.