

Datasheet for ABIN6244252  
**anti-DEFB107A antibody (C-Term)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	400 µL
Target:	DEFB107A
Binding Specificity:	AA 56-90, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DEFB107A antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Western Blotting (WB)

## Product Details

Immunogen:	This DEFB107A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 56-90 amino acids from the C-terminal region of human DEFB107A.
Clone:	RB47863
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	DEFB107A
Alternative Name:	DEFB107A ( <a href="#">DEFB107A Products</a> )
Background:	Has antibacterial activity (Potential).

## Target Details

Molecular Weight: 7846

UniProt: [Q8IZN7](#)

## Application Details

Application Notes: WB: 1:1000. IHC-P: 1:25

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

## Images



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemical analysis of paraffin-embedded H. testis section using DEFB107A Antibody (C-term) (ABIN6244252 and ABIN6577438). (ABIN6244252 and ABIN6577438) was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



### Western Blotting

**Image 2.** Western blot analysis of lysate from human testis tissue lysate, using DEFB107A Antibody (C-term) (ABIN6244252 and ABIN6577438). (ABIN6244252 and ABIN6577438) was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 µg.