antibodies - online.com







anti-TRBC1 antibody (AA 104-136)

Images



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

0.101.101.		
Quantity:	400 μL	
Target:	TRBC1	
Binding Specificity:	AA 104-136	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TRBC1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This TRBC1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 104-136 amino acids from the Central region of human TRBC1.	
Clone:	RB45122	
Isotype:	Ig Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	TRBC1	
Alternative Name:	TRBC1 (TRBC1 Products)	
Molecular Weight:	19769	

Target Details

UniProt: P01850

Application Details


Application Notes:	WB: 1:500. WB: 1:500. WB: 1:1000

Restrictions: For Research Use only

Handling


Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	


Images



Western Blotting

Image 1. Western blot analysis of lysate from cell line, using TRBC1 Antibody (Center) (ABIN6244260 and ABIN6577681). (ABIN6244260 and ABIN6577681) was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20 μ g.





Western Blotting

Image 2. All lanes: Anti-TRBC1 Antibody (Center) at 1:500 dilution Lane 1: Molt-4 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size: 34 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Western Blotting

Image 3. All lanes: Anti-TRBC1 Antibody (Center) at 1:500 dilution Lane 1: Molt-4 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size: 36 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.