antibodies .- online.com







anti-RAD9B antibody (AA 394-428)







\sim	
()\/\	rview
\cup	1 410 44

Quantity:	400 μL
Target:	RAD9B
Binding Specificity:	AA 394-428
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAD9B antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Target/Specificity: This RAD9B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of
Immunogen:	
Immunogen: Clone:	conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of
	conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human RAD9B.
Clone:	conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human RAD9B. RB49841
Clone:	conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human RAD9B. RB49841 Ig Fraction
Clone: Isotype: Purification:	conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human RAD9B. RB49841 Ig Fraction

Target Details


Background:	Cell cycle checkpoint control protein RAD9B, DNA repair exonuclease rad9 homolog B, hRAD9B, RAD9B
Molecular Weight:	47832
UniProt:	Q6WBX8

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	

Images

Format:



Liquid

Western Blotting

Image 1. Western blot analysis of lysates from THP-1, U-251 MG, HL-60 cell line (from left to right), using RAD9B Antibody (C-term) (ABIN6244327 and ABIN6577414). (ABIN6244327 and ABIN6577414) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 μg per lane.