

Datasheet for ABIN6252976

anti-ITGAL antibody





Overview

| Quantity: | 100 μg |
|--------------|--------------------------------------|
| Target: | ITGAL |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This ITGAL antibody is un-conjugated |
| Application: | Flow Cytometry (FACS) |

Product Details

| Purpose: | CD11a Unconjugated Antibody |
|------------------|--|
| Immunogen: | Human HLA-DRCTL line cytolytic Tlymphocytes |
| Clone: | TS1-22 |
| Isotype: | IgG1, kappa |
| Characteristics: | The clone recognizes a 170-180 kD alpha chain of a type I transmembrane glycoprotein also known as CD11a or LFA-1a. It is also known as LFA-1 α . CD11a non-covalently associates with integrin β 2 (CD18) to form LFA-1. CD11a is abundantly expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils but does not express by the non-hematopoietic tissues and platelets. Up-regulation of CD11a on activated lymphocytes is required to induce their immune response. Absence or reduced CD11a expression on tumor cells helps to escape from regular immune surveillance and it may be one of reason of in vivo tumor generation. |

Product Details

| Purification: | Purified |
|---------------|-----------|
| Purity: | >95 % |
| Grade: | GMP Grade |

Target Details


| Target: | ITGAL |
|-------------------|--|
| Alternative Name: | CD11a (ITGAL Products) |
| UniProt: | P20701 |
| Pathways: | Activated T Cell Proliferation, Integrin Complex |

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Buffer: | PBS pH 7.2, 0.1 % (w/v) BSA, 0.09 % (w/v) sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |



Flow Cytometry

Image 1.