

Datasheet for ABIN6253114

anti-CD4 antibody (iFluor™700)





Go to Product page

Overview

Quantity:	100 tests
Target:	CD4
Reactivity:	Human, Rhesus Monkey, Cynomolgus
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD4 antibody is conjugated to iFluor™700
Application:	Flow Cytometry (FACS)


Product Details


Purpose:	CD4 iFluor™ 700 Antibody
Immunogen:	Human peripheral blood T lymphocytes
Clone:	OKT4
Isotype:	IgG2b, kappa
Characteristics:	CD4 is a 55kD type I transmembrane glycoprotein and a member of the immunoglobulin superfamily. It is a specific marker for T lymphocytes, monocytes, macrophages, and dendritic cells. Through interaction of MHC-II, CD4 facilitates cell-cell interaction, thymic differentiation, and activation of downstream signaling cascades. HIV infection of T-cells is instigated through binding of HIV to CD4. The OKT4 binds to the D3 domain of CD4 and does not inhibit HIV binding.
Purification:	Purified
Purity:	>95 %

Product Details GMP Grade Grade: **Target Details** CD4 Target: Alternative Name CD4 (CD4 Products) NCBI Accession: NM_000616 UniProt: P01730 Pathways: TCR Signaling, Maintenance of Protein Location, CXCR4-mediated Signaling Events **Application Details** Optimal working dilution should be determined by the investigator. Application Notes: Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS pH 7.2, 0.2 % (w/v) BSA, 0.09 % (w/v) sodium azide Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. 4°C Storage:

2-8°C, Conjugated antibodies should never be frozen.

Storage Comment:





Flow Cytometry

Image 1.

Image 2. Lymphocytes gated PBMC stained with iFluor700 conjugated anti-human CD4 (clone OKT 4, red histogram). Lymphocytes gated PBMC with FMO control (blue histogram). The dara were generated in Beckman Coulter CvtoFLEX S Flow Cvtometer and analyzed by using FlowJo software.