

Datasheet for ABIN6253220

CD137 Protein (AA 19-184) (Fc Tag)



Go to Product page

| () | ve | r\/i | Δ | ۱۸/ |
|---------|-----|------|---|-----|
| \circ | V C | 1 V | | v v |

| Quantity: | 100 μg |
|-------------------------------|---|
| Target: | CD137 (TNFRSF9) |
| Protein Characteristics: | AA 19-184 |
| Origin: | Mouse |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CD137 protein is labelled with Fc Tag. |

Product Details

| Purpose: | CD137 (human):Fc (mouse) (rec.) |
|------------------|--|
| Specificity: | The extracellular domain of human CD137 [4-1BB] (aa 19-184) is fused to the N-terminus of the Fc region of mouse IgG2a. |
| Characteristics: | Protein. The extracellular domain of human CD137 [4-1BB] (aa 19-184) is fused to the N-terminus of the Fc region of mouse IgG2a. Source: HEK 293 cells. Endotoxin content: <5EU/mg protein (LAL test, Lonza). Lyophilized from 0.2µm-filtered solution in PBS. Purity: >98 % (SDS-PAGE). Human CD137 (4-1BB) is a costimulatory molecule of the tumor necrosis factor (TNF) receptor superfamily. The glycoprotein 4-1BB is expressed mainly on activated CD4+ and CD8+ T cells and binds to a high-affinity ligand (4-1BBL) expressed on several antigen-presenting cells such as macrophages and activated B cells. Upon ligand binding, 4-1BB is associated with the tumor receptor-associated factors (TRAF), the adaptor protein and mediates downstream signaling events including the activation of NF-kappaB and cytokine production. 4-1BB signaling either by binding to 4-1BBL or by antibody ligation delivers signals for T cell activation and |

growth as well as monocyte proliferation and B cell survival, and plays a important role in the amplification of T cell-mediated immune responses. In contrast, it can also enhance activation-induced T cell apoptosis when triggered by engagement of the TCR/CD3 complex. In addition, the 4-1BB/4-1BBL costimulatory pathway has been shown to augment secondary CTL responses to several viruses and increase antitumor immunity. 4-1BB is therefore a promising candidate for immunotherapy of human cancer.

Purity:

>98 % (SDS-PAGE)

Endotoxin Level:

<5EU/mg protein (LAL test, Lonza).

Biological Activity Comment:

Measured by its binding ability in a functional ELISA.

Target Details

Target:

CD137 (TNFRSF9)

Alternative Name:

CD137 (TNFRSF9 Products)

Background:

Alternate Names/Synonyms: Tumor Necrosis Factor Receptor Superfamily Member 9, TNFRSF9, 4-1BB, 4-1BB Ligand Receptor T Cell, Antigen 4-1BB Homolog, T Cell Antigen ILA, CD137 Antigen, CDw137, ILA, MGC2172

Product Description: Human CD137 (4-1BB) is a costimulatory molecule of the tumor necrosis factor (TNF) receptor superfamily. The glycoprotein 4-1BB is expressed mainly on activated CD4+ and CD8+ T cells and binds to a high-affinity ligand (4-1BBL) expressed on several antigen-presenting cells such as macrophages and activated B cells. Upon ligand binding, 4-1BB is associated with the tumor receptor-associated factors (TRAF), the adaptor protein and mediates downstream signaling events including the activation of NF-kappaB and cytokine production. 4-1BB signaling either by binding to 4-1BBL or by antibody ligation delivers signals for T cell activation and growth as well as monocyte proliferation and B cell survival, and plays a important role in the amplification of T cell-mediated immune responses. In contrast, it can also enhance activation-induced T cell apoptosis when triggered by engagement of the TCR/CD3 complex. In addition, the 4-1BB/4-1BBL costimulatory pathway has been shown to augment secondary CTL responses to several viruses and increase antitumor immunity. 4-1BB is therefore a promising candidate for immunotherapy of human cancer.

Molecular Weight:

~55kDa (SDS-PAGE)

Pathways:

Cancer Immune Checkpoints

Application Details

| Restrictions: | For Research Use only |
|------------------|--|
| Handling | |
| Format: | Lyophilized |
| Concentration: | Lot specific |
| Buffer: | Lyophilized from 0.2µm-filtered solution in PBS. |
| Handling Advice: | Avoid freeze/thaw cycles. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Short Term Storage: +4°C |
| | Long Term Storage: -20°C |
| | Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots |

are stable for up to 3 months when stored at -20°C.