



[Go to Product page](#)

Datasheet for ABIN6253402
HMGB1 Protein (AA 24-215) (His tag)

Overview

Quantity:	25 µg
Target:	HMGB1
Protein Characteristics:	AA 24-215
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This HMGB1 protein is labelled with His tag.

Product Details

Purpose:	HMGB1 (human) (rec.) (His)
Specificity:	Human HMGB1 (aa 24-215) is fused at the N-terminus to a His-tag.
Characteristics:	<p>Protein. Human HMGB1 (aa 24-215) is fused at the N-terminus to a His-tag. Source: E. coli.</p> <p>Endotoxin content: <0.1EU/µg protein (LAL test, Lonza). Lyophilized from a concentrated sterile solution containing 50 mM Tris-Hcl buffer (pH 8.0) and 100 mM NaCl. Purity: >90 % (SDS-PAGE). HMGB1 was originally discovered as an essential DNA-binding protein for regulating p53, NF-kappaB and other important proteins. It is secreted from activated dendritic cells, macrophage and necrotic cells, and acts as a ligand for RAGE, TLR-2 and TLR-4 expressed on surrounding cells. As a result, HMGB1 activates Rac, CDC42 and NF-kappaB inducing localized innate immunity of damaged tissue, tissue regeneration by recruitment of stem cells and hemostasis by induction of tissue factor expression. HMGB1 is also causative agent of various diseases as it causes localized inflammation such as arteriosclerosis, chronic rheumatoid arthritis and nephritis.</p>

Product Details

Purity:	>90 % (SDS-PAGE)
Endotoxin Level:	<0.1EU/μg protein (LAL test, Lonza).

Target Details

Target:	HMGB1
Alternative Name:	HMGB1 (HMGB1 Products)
Background:	<p>Alternate Names/Synonyms: High Mobility Group Protein B1</p> <p>Product Description: HMGB1 was originally discovered as an essential DNA-binding protein for regulating p53, NF-kappaB and other important proteins. It is secreted from activated dendritic cells, macrophage and necrotic cells, and acts as a ligand for RAGE, TLR-2 and TLR-4 expressed on surrounding cells. As a result, HMGB1 activates Rac, CDC42 and NF-kappaB inducing localized innate immunity of damaged tissue, tissue regeneration by recruitment of stem cells and hemostasis by induction of tissue factor expression. HMGB1 is also causative agent of various diseases as it causes localized inflammation such as arteriosclerosis, chronic rheumatoid arthritis and nephritis.</p>
NCBI Accession:	NP_002119
Pathways:	p53 Signaling , Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development , Positive Regulation of Endopeptidase Activity , Regulation of Carbohydrate Metabolic Process , Toll-Like Receptors Cascades , Smooth Muscle Cell Migration , Inflammasome

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized from a concentrated sterile solution containing 50 mM Tris-Hcl buffer (pH 8.0) and 100 mM NaCl.
Handling Advice:	Avoid freeze/thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Short Term Storage: +4°C

Handling

Long Term Storage: -20°C

Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.