

[Go to Product page](#)

Datasheet for ABIN6253411

TIMD4 Protein (AA 25-315) (Fc Tag)

Overview

Quantity:	100 µg
Target:	TIMD4
Protein Characteristics:	AA 25-315
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TIMD4 protein is labelled with Fc Tag.

Product Details

Purpose:	Tim-4 (human):Fc (mouse) (rec.)
Specificity:	The extracellular domain of human Tim-4 (aa 25-315) is fused to the N-terminus of the Fc region of mouse IgG2a.
Characteristics:	Protein. The extracellular domain of human Tim-4 (aa 25-315) is fused to the N-terminus of the Fc region of mouse IgG2a. Source: HEK 293 cells. Endotoxin content: <5EU/mg protein (LAL test, Lonza). Lyophilized from 0.2µm-filtered solution in PBS. Purity: >98 % (SDS-PAGE). The TIM (T cell/transmembrane, immunoglobulin and mucin) family plays a critical role in regulating immune responses, including allergy, asthma, transplant tolerance, autoimmunity and the response to viral infections. The unique structure of TIM immunoglobulin variable region domains allows highly specific recognition of phosphatidylserine (PtdSer), exposed on the surface of apoptotic cells. TIM-4 (T cell, immunoglobulin, Mucin-4), also known as SMUCKLER, is a 60 kDa member of the TIM family of immune regulating proteins. TIM-4 is exclusively expressed on antigen-presenting cells, where it mediates phagocytosis of apoptotic cells and

Product Details

plays an important role in maintaining tolerance. TIM-4 binds specifically to TIM-1 which is also the cellular receptor for the hepatitis A virus, and has been implicated in the development of asthma. Among hematopoietic cells, TIM-1 is expressed on activated B and T cells, preferentially in the Th2 subset of CD4+ T cells. The interaction of TIM-4 with TIM-1 induces costimulatory and hyperproliferative signals in T cells.

Purity: >98 % (SDS-PAGE)

Endotoxin Level: <5EU/mg protein (LAL test, Lonza).

Biological Activity Comment: Measured by its ability to inhibit anti-CD3-induced proliferation of stimulated human T cells.

Target Details

Target: TIMD4

Alternative Name: Tim-4 ([TIMD4 Products](#))

Background: Alternate Names/Synonyms: TIM4, TIMD4, T Cell Immunoglobulin and Mucin Domain-containing Protein 4, T Cell Membrane Protein 4

Product Description: The TIM (T cell/transmembrane, immunoglobulin and mucin) family plays a critical role in regulating immune responses, including allergy, asthma, transplant tolerance, autoimmunity and the response to viral infections. The unique structure of TIM immunoglobulin variable region domains allows highly specific recognition of phosphatidylserine (PtdSer), exposed on the surface of apoptotic cells. TIM-4 (T cell, immunoglobulin, Mucin-4), also known as SMUCKLER, is a 60 kDa member of the TIM family of immune regulating proteins. TIM-4 is exclusively expressed on antigen-presenting cells, where it mediates phagocytosis of apoptotic cells and plays an important role in maintaining tolerance. TIM-4 binds specifically to TIM-1 which is also the cellular receptor for the hepatitis A virus, and has been implicated in the development of asthma. Among hematopoietic cells, TIM-1 is expressed on activated B and T cells, preferentially in the Th2 subset of CD4+ T cells. The interaction of TIM-4 with TIM-1 induces costimulatory and hyperproliferative signals in T cells.

Molecular Weight: ~80kDa (SDS-PAGE)

NCBI Accession: [NP_612388](#)

Pathways: [Cancer Immune Checkpoints](#)

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized from 0.2µm-filtered solution in PBS.
Handling Advice:	Avoid freeze/thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.