antibodies.com

# Datasheet for ABIN6253546 IGF1R Protein (AA 31-932) (His tag)

2 Images



#### Overview

Quantity:	100 µg
Target:	IGF1R
Protein Characteristics:	AA 31-932
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IGF1R protein is labelled with His tag.

## Product Details


Sequence:	AA 31-932
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.


### Target Details

Target:	IGF1R
Alternative Name:	IGF-I R (IGF1R Products)
Background:	The Insulin-like Growth Factor 1 Receptor (IGF1) is also known as CD221, JTK13. and is a
	transmembrane receptor that is activated by IGF-1 and by the related growth factor IGF-2. It
	belongs to the large class of tyrosine kinase receptors. This receptor mediates the effects of
	IGF-1, which is a polypeptide protein hormone similar in molecular structure to insulin. IGF1R is
	make up of two alpha subunits and two beta subunits ,the Both the $lpha$ and $eta$ subunits are

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6253546 | 01/16/2024 | Copyright antibodies-online. All rights reserved.

	synthesized from a single mRNA precursor. The precursor is then glycosylated, proteolytically
	cleaved, and crosslinked by cysteine bonds to form a functional transmembrane $lphaeta$ chain.The c
	chains are located extracellularly while the $\beta$ subunit spans the membrane and are responsible
	for intracellular signal transduction upon ligand stimulation. IGF1R have a binding site for ATP,
	which is used to provide the phosphates for autophosphorylation. There is a 60 $\%$ homology
	between IGF1R and the insulin receptor. In response to ligand binding, the $lpha$ chains induce the
	tyrosine autophosphorylation of the $eta$ chains. This event triggers a cascade of intracellular
	signaling that, while somewhat cell type specific, often promotes cell survival and cell
	proliferation.
Molecular Weight:	81.1 kDa, 23.8 kDa
NCBI Accession:	XP_005560632
Pathways:	RTK Signaling, Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic
	Process, Autophagy
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C





#### ELISA

**Image 1.** Immobilized Cynomolgus IGF-I R, His Tag (ABIN6253205,ABIN6253546) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Human IGF-I, Fc Tag (ABIN2181263,ABIN2181262) with a linear range of 0.02-0.3  $\mu$ g/mL (QC tested).

#### SDS-PAGE

**Image 2.** Cynomolgus IGF-I R, His Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 % .

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6253546 | 01/16/2024 | Copyright antibodies-online. All rights reserved.