

Datasheet for ABIN6253646

BAFF Protein (AA 127-309) (Fc Tag)**3** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	BAFF (TNFSF13B)
Protein Characteristics:	AA 127-309
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This BAFF protein is labelled with Fc Tag.

Product Details

Sequence:	AA 127-309
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 0.1 EU per µg by the LAL method.

Target Details

Target:	BAFF (TNFSF13B)
Alternative Name:	BAFF (TNFSF13B Products)
Background:	B-cell activating factor (BAFF) is also known as tumor necrosis factor ligand superfamily member 13B , TNFSF13B, BAFF, B Lymphocyte Stimulator (BLyS) , cluster of differentiation 257 (CD257), DTL, TNF- and APOL-related leukocyte expressed ligand (TALL-1), THANK, TNFSF20, ZTNF4, and is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This

Target Details

cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, and acts as a potent B cell activator. It has been also shown to play an important role in the proliferation and differentiation of B cells. It is expressed as transmembrane protein on various cell types including monocytes, dendritic cells and bone marrow stromal cells. BAFF is the natural ligand of three unusual tumor necrosis factor receptors named BAFF-R, TACI, and BCMA, all of which have differing binding affinities for it. These receptors are expressed mainly on mature B lymphocytes (TACI is also found on a subset of T-cells and BCMA on plasma cells). TACI binds worst since its affinity is higher for a protein similar to BAFF, called a proliferation-inducing ligand (APRIL). BCMA displays an intermediate binding phenotype and will work with either BAFF or APRIL to varying degrees. Signaling through BAFF-R and BCMA stimulates B lymphocytes to undergo proliferation and to counter apoptosis. All these ligands act as heterotrimers (i.e. three of the same molecule) interacting with heterotrimeric receptors, although BAFF has been known to be active as either a hetero- or homotrimer. BAFF acts as a potent B cell activator and has been shown to play an important role in the proliferation and differentiation of B cells.




Molecular Weight:	47.7 kDa
Pathways:	NF-kappaB Signaling , Production of Molecular Mediator of Immune Response

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Buffer:	50 mM Tris, 100 mM Glycine, pH 7.5
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C



ELISA

Image 1. Immobilized Mouse BAFF, Mouse IgG2a Fc Tag, low endotoxin (ABIN5954906,ABIN6253646) at 5 µg/mL (100 µL/well) can bind Mouse BAFFR, Fc Tag (ABIN5526590,ABIN5526591) with a linear range of 0.4-3 ng/mL (Routinely tested).

SDS-PAGE

Image 2. Mouse BAFF, Mouse IgG2a Fc Tag, low endotoxin on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .

ELISA

Image 3. Immobilized Mouse BAFF, Mouse IgG2a Fc Tag (ABIN5954906,ABIN6253646) at 5 µg/mL (100 µL/well) can bind Mouse BCMA, Fc Tag (ABIN5674646,ABIN6253667) with a linear range of 2-10 ng/mL (QC tested).