

Datasheet for ABIN625421

MME ELISA Kit





Overview

Quantity:	96 tests
Target:	MME
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	61-15000 pg/mL
Minimum Detection Limit:	61 pg/mL
Application:	ELISA

Product Details

Product Details	
Purpose:	Mouse Neprilysin ELISA Kit for Cell Culture Supernatants, Plasma, and Serum samples.
Sample Type:	Cell Culture Supernatant, Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes mouse Neprilysin.
Sensitivity:	61 pg/mL
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data
Components:	Pre-Coated 96-well Strip Microplate

Product Details

- · Wash Buffer
- · Stop Solution
- · Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- · TMB One-Step Substrate

Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μL to 1 μL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 μL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:	MME
Alternative Name:	Neprilysin (MME Products)
Target Type:	Chemical
Background:	Gene Names: Mme Protein names: Neprilysin (EC 3.4.24.11) (Atriopeptidase) (Enkephalinase) (Neutral endopeptidase 24.11) (NEP) (Neutral endopeptidase) (Skin fibroblast elastase) (SFE) (CD antigen CD10)
Gene ID:	17380
UniProt:	Q61391
Pathways:	RTK Signaling, Peptide Hormone Metabolism, Regulation of Systemic Arterial Blood Pressure by Hormones, Smooth Muscle Cell Migration

Application Details

Application Notes:	Recommended Dilution for serum and plasma samples2 fold
Sample Volume:	100 μL
Plate:	Pre-coated Pre-coated

Application Details

Protocol:

- 1. Prepare all reagents, samples and standards as instructed in the manual.
- 2. Add 100 µL of standard or sample to each well.
- 3. Incubate 2.5 h at RT or O/N at 4 °C.
- 4. Add 100 μL of prepared biotin antibody to each well.
- 5. Incubate 1 h at RT.
- 6. Add 100 µL of prepared Streptavidin solution to each well.
- 7. Incubate 45 min at RT.
- 8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions:

For Research Use only

Handling

Stora	age:
-------	------

-20 °C

Storage Comment:


The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4° C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date:

6 months

Images





Mouse Neprilysin concentration (pg/ml)

ELISA

Image 1.