antibodies - online.com







anti-Chaperonin GroEL (GroEL) (C-Term) antibody



Image




| \sim | | | |
|--------|----------------------------|------|-----------|
| | $ \backslash / \square $ | r\/I | \square |

| Quantity: | 600 µg | |
|----------------------|---|--|
| Target: | Chaperonin GroEL (GroEL) | |
| Binding Specificity: | C-Term | |
| Reactivity: | Burkholderia, Pseudomonas | |
| Host: | Goat | |
| Clonality: | Polyclonal | |
| Conjugate: | Un-conjugated | |
| Application: | Western Blotting (WB) | |
| Product Details | | |
| Immunogen: | Purified recombinant peptide derived from within residues 400 aa to the C-terminus of | |
| | Burkholderia GroEL produced in E. coli. | |
| Isotype: | IgG | |
| Specificity: | The Ab recognizes the endogenous protein in Burkholderia and Pseudomonas by Western blot. | |
| Characteristics: | Chaperonin GroEL protein | |
| Purification: | epitope affinity purified | |
| Target Details | | |
| Target: | Chaperonin GroEL (GroEL) | |
| Alternative Name: | GroEL (GroEL Products) | |

Application Details

| Application Notes: Optimal working dilution should be determined by the investigator. | | |
|---|--|--|
| Restrictions: | For Research Use only | |
| Handling | | |
| Handling Advice: | The antibody solution should be gently mixed before use. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | For continuous use, store at 2-8 deg,C for one-two days. For extended storage, store in -20 deg,C freezer. Working dilution samples should be discarded if not used within 12 hours. | |

Images



Western Blotting

Image 1. Anti-GroEL Ab at 1/2,500 dilution, $50-100~\mu g$ of total protein per lane, rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution,