antibodies -online.com





anti-FTCD antibody (N-Term)



Images



Go to Product page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

0.01.11011		
Quantity:	600 µg	
Target:	FTCD	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat, Pig, Dog, Rabbit, Horse, Guinea Pig, Cow, Monkey, Avian, Cat, Chicken,	
	Donkey, Goat, Hamster, Sheep	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This FTCD antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF)	
Product Details		
Immunogen:	Purified recombinant peptide derived from within residues 50 aa to the N-terminus of human	
	FTCD produced in E. coli.	
Isotype:	IgG	
Specificity:	Detects a band of 55-60 kDa by Western blot in the following canine, human, monkey, mouse,	
	rat whole cell lysates.	
Characteristics:	Formimidoyltransferase cyclodeaminase	
Purification:	epitope affinity purified	
Target Details		
Target:	FTCD	

Target Details

Alternative Name:	FTCD (FTCD Products)
-------------------	----------------------


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Handling Advice:	The antibody solution should be gently mixed before use.
Storage:	4 °C,-20 °C
Storage Comment:	For continuous use, store at 2-8 deg,C for one-two days. For extended storage, store in -20
	deg,C freezer. Working dilution samples should be discarded if not used within 12 hours.

Images




Immunohistochemistry

Image 1.



Immunofluorescence

Image 2. Immunofluorescence - anti-FTCD Ab at 1/100 dilution in NHI/3T3 cells were fixed with methanol and permeabilized With 0.1% saponin,



Western Blotting

Image 3. Anti-FTCO Ab at 1/1,000 dilution,at 100 lane, Rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution,

Please check the product details page for more images. Overall 9 images are available for ABIN6254252.