antibodies -online.com





anti-FADD antibody (pSer194)

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	FADD
Binding Specificity:	pSer194
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FADD antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details


Immunogen:	A synthesized peptide derived from human FADD around the phosphorylation site of Ser194.
Isotype:	IgG
Specificity:	Phospho-FADD (Ser194) Antibody detects endogenous levels of FADD only when phosphorylated at Ser194.
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

Target:	FADD
Alternative Name:	FADD (FADD Products)


Target Details

rarget betails	
Background:	Description: Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling. Gene: FADD
Molecular Weight:	23kDa
Gene ID:	8772
UniProt:	Q13158
Pathways:	Apoptosis, TLR Signaling, Activation of Innate immune Response, Positive Regulation of Endopeptidase Activity, Toll-Like Receptors Cascades
Application Details	
Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of Phospho-FADD (Ser194) using COLO205 whole cell lysates



Immunofluorescence (fixed cells)

Image 2. ABIN6274076 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.