antibodies -online.com





anti-IKBKG antibody (pSer31)

Images



\sim					
	1//	⊃r	V/I	Φ\	Λ

Quantity:	100 μL	
Target:	IKBKG	
Binding Specificity:	pSer31	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IKBKG antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	A synthesized peptide derived from human IKK gamma around the phosphorylation site of Ser31.	
Isotype:	IgG	
Specificity:	Phospho-IKK gamma (Ser31) Antibody detects endogenous levels of IKK gamma only when phosphorylated at Serine 31.	
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog	
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential	

chromatography on phospho- and non-phospho-peptide affinity columns.


Target Details

Target:	IKBKG	
Alternative Name:	IKBKG (IKBKG Products)	
Background:	Description: Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-	
	kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the	
	degradation of the inhibitor. Its binding to scaffolding polyubiquitin seems to play a role in IKK	
	activation by multiple signaling receptor pathways. However, the specific type of polyubiquitin	
	recognized upon cell stimulation (either 'Lys-63'-linked or linear polyubiquitin) and its functional	
	importance is reported conflictingly. Also considered to be a mediator for TAX activation of NF-	
	kappa-B. Could be implicated in NF-kappa-B-mediated protection from cytokine toxicity.	
	Essential for viral activation of IRF3. Involved in TLR3- and IFIH1-mediated antiviral innate response, this function requires 'Lys-27'-linked polyubiquitination.	
	Gene: IKBKG	
Molocular Waight		
Molecular Weight:	48kDa	
Gene ID:	8517 	
UniProt:	Q9Y6K9	
Pathways:	NF-kappaB Signaling, RTK Signaling, TCR Signaling, TLR Signaling, Fc-epsilon Receptor	
	Signaling Pathway, Activation of Innate immune Response, M Phase, Production of Molecular	
	Mediator of Immune Response, Hepatitis C, Protein targeting to Nucleus, Toll-Like Receptors	
	Cascades, BCR Signaling, Ubiquitin Proteasome Pathway, S100 Proteins	
Application Details		
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	

Handling


	should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	

Images




Immunofluorescence (fixed cells)

Image 1. ABIN6267704 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 2. Western blot analysis of IKK- gamma phosphorylation expression in TNF-a treated 293 whole cell lysates, The lane on the left is treated with the antigenspecific peptide.



Immunohistochemistry

Image 3. ABIN6267704 at 1/200 staining human testis tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.