antibodies -online.com





anti-Tyrosine Hydroxylase antibody (pSer40)



Images



Overview	
Quantity:	

Quantity:	100 μL
Target:	Tyrosine Hydroxylase (TH)
Binding Specificity:	pSer40
Reactivity:	Rat, Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tyrosine Hydroxylase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), ELISA, Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human Tyrosine Hydroxylase around the phosphorylation site of Serine 40
Isotype:	IgG
Specificity:	Phospho-Tyrosine Hydroxylase (Ser40) Antibody detects endogenous levels of Tyrosine Hydroxylase only when phosphorylated at Serine 40
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details


Target:	Tyrosine Hydroxylase (TH)
Alternative Name:	Tyrosine Hydroxylase (TH Products)
Background:	Description: Plays an important role in the physiology of adrenergic neurons. Gene: TH
Molecular Weight:	60kDa
Gene ID:	7054
UniProt:	P07101
Pathways:	Dopaminergic Neurogenesis, Response to Water Deprivation, Sensory Perception of Sound, Carbohydrate Homeostasis, Feeding Behaviour

Application Details

Application Notes:	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months




Immunohistochemistry

Image 1. ABIN6267326 at 1/200 staining Rat brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunofluorescence (fixed cells)

Image 2. ABIN6267326 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 3. Western blot analysis of Tyrosine Hydroxylase phosphorylation expression in RAW264.7 whole cell lysates,The lane on the left is treated with the antigenspecific peptide.

Please check the product details page for more images. Overall 5 images are available for ABIN6255304.