

Datasheet for ABIN6255425

anti-ENOS antibody (pSer615)





\sim	+ ~	Produ	10+	0000
(¬()	1()	PICKII	10,1	11200

Overview

Quantity:	100 μL
Target:	ENOS (NOS3)
Binding Specificity:	pSer615
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ENOS antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human eNOS around the phosphorylation site of Ser615.
Isotype:	IgG
Specificity:	Phospho-eNOS (Ser615) Antibody detects endogenous levels of eNOS only when phosphorylated at Sersine 615.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Target Details	
Target:	ENOS (NOS3)


Target Details


Alternative Name:	NOS3 (NOS3 Products)
Background:	Description: Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets. Gene: NOS3
Molecular Weight:	140kDa
Gene ID:	4846
Gene ID: UniProt:	4846 P29474

Application Details

Application Notes:	WB 1:500-1:2000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6266457 staining K562 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis on K562 cell lysate using Phospho-eNOS(Ser615) Antibody. The lane on the left is treated with the antigen-specific peptide.