antibodies -online.com





anti-HNRNPD/AUF1 antibody (pSer83)

4 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	HNRNPD/AUF1 (HNRNPD)
Binding Specificity:	pSer83
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPD/AUF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human hnRPD around the phosphorylation site of Ser83.
Isotype:	IgG
Specificity:	Phospho-hnRPD (Ser83) Antibody detects endogenous levels of hnRPD only when phosphorylated at Ser83.
Predicted Reactivity:	Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

Target: HNRNPD/AUF1 (HNRNPD)


Target Details


Alternative Name:	HNRNPD (HNRNPD Products)
Background:	Description: Binds with high affinity to RNA molecules that contain AU-rich elements (AREs)
	found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Also binds to double-
	and single-stranded DNA sequences in a specific manner and functions a transcription factor.
	Each of the RNA-binding domains specifically can bind solely to a single-stranded non-
	monotonous 5'-UUAG-3' sequence and also weaker to the single-stranded 5'-TTAGGG-3'
	telomeric DNA repeat. Binds RNA oligonucleotides with 5'-UUAGGG-3' repeats more tightly than
	the telomeric single-stranded DNA 5'-TTAGGG-3' repeats. Binding of RRM1 to DNA inhibits the
	formation of DNA quadruplex structure which may play a role in telomere elongation. May be
	involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins
	in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated
	by the major coding-region determinant of instability (mCRD) domain. May play a role in the
	regulation of the rhythmic expression of circadian clock core genes. Directly binds to the 3'UTR
	of CRY1 mRNA and induces CRY1 rhythmic translation. May also be involved in the regulation
	of PER2 translation.
	Gene: HNRNPD
Molecular Weight:	38 kDa
Gene ID:	3184
UniProt:	Q14103
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.


Handling

Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images







Immunofluorescence (fixed cells)

Image 1. ABIN6268985 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Western Blotting

Image 2. Western blot analysis of hnRPD (Phospho-Ser83) expression in 293 cell extract. The lane on the left is treated with the antigen-specific peptide.

Immunofluorescence (fixed cells)

Image 3. ABIN6268985 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25;ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37;ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Please check the product details page for more images. Overall 4 images are available for ABIN6255528.