antibodies -online.com





anti-RPS6KB1 antibody (pThr229)



Images



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	RPS6KB1
Binding Specificity:	pThr229
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RPS6KB1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human p70 S6 Kinase around the phosphorylation site of Thr229.	
Isotype:	IgG	
Specificity:	Phospho-p70 S6 Kinase (Thr229) Antibody detects endogenous levels of p70 S6 Kinase only when phosphorylated at Thr252, which site historically referenced as Thr229.	
Predicted Reactivity:	Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus	
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	

Target Details RPS6KB1 Target: Alternative Name: RPS6KB1 (RPS6KB1 Products) Background: Description: Serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4, targeting it for ubiquitination and subsequent proteolysis. Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR. In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2. Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling. Mediates cell survival by phosphorylating the pro-apoptotic protein BAD and suppressing its proapoptotic function. Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex. The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function. Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1. In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B. May be involved in cytoskeletal rearrangement through binding to neurabin. Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR (PubMed:11500364, PubMed:12801526, PubMed:14673156, PubMed:15071500, PubMed:15341740, PubMed:16286006, PubMed:17052453, PubMed:17053147, PubMed:17936702, PubMed:18952604, PubMed:19085255, PubMed:19720745, PubMed:19935711, PubMed:19995915, PubMed:23429703). Following activation by mTORC1, phosphorylates EPRS and thereby plays a key role in fatty acid uptake by adipocytes and also most probably in interferon-gammainduced translation inhibition (PubMed:28178239).

Molecular Weight:

70kDa

Gene: RPS6KB1

Target Details


Gene ID:	6198
UniProt:	P23443
Pathways:	PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, Regulation of Cell Size, Skeletal Muscle Fiber Development, Feeding Behaviour, G-protein mediated Events, Smooth Muscle Cell
	Migration, Interaction of EGFR with phospholipase C-gamma, Warburg Effect

Application Details

Expiry Date:


Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	

12 months




Western Blotting

Image 1. Western blot analysis of p70 S6 Kinase phosphorylation expression in Jurkat whole cell lysates,The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 2. ABIN6267437 at 1/200 staining Mouse heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunohistochemistry

Image 3. ABIN6267437 at 1/200 staining Rat lung tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 6 images are available for ABIN6255636.