antibodies -online.com





Datasheet for ABIN6255994

anti-KIT antibody (pTyr703)

Images



\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Overview	
Quantity:	100 μL
Target:	KIT
Binding Specificity:	pTyr703
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIT antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogon:	A synthesized poptide derived from human c-Kit around the phospherylation site of Tyr703

Immunogen:	A synthesized peptide derived from human c-Kit around the phosphorylation site of Tyr703.
Isotype:	IgG
Specificity:	Phospho-c-Kit (Tyr703) Antibody detects endogenous levels of c-Kit only when phosphorylated at Tyrosine 703.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog,Chicken
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

Target: KIT


Target Details

Alternative Name:	KIT (KIT Products)		
Background:	Description: Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine		
	KITLG/SCF and plays an essential role in the regulation of cell survival and proliferation,		
	hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and		
	function, and in melanogenesis. In response to KITLG/SCF binding, KIT can activate several		
	signaling pathways. Phosphorylates PIK3R1, PLCG1, SH2B2/APS and CBL. Activates the AKT1		
	signaling pathway by phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol		
	3-kinase. Activated KIT also transmits signals via GRB2 and activation of RAS, RAF1 and the		
	MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family		
	members STAT1, STAT3, STAT5A and STAT5B. Activation of PLCG1 leads to the production of		
	the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KIT signaling is		
	modulated by protein phosphatases, and by rapid internalization and degradation of the		
	receptor. Activated KIT promotes phosphorylation of the protein phosphatases PTPN6/SHP-1		
	and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B. Promotes		
	phosphorylation of PIK3R1, CBL, CRK (isoform Crk-II), LYN, MAPK1/ERK2 and/or MAPK3/ERK1		
	PLCG1, SRC and SHC1.		
	Gene: KIT		
Molecular Weight:	145kDa		
Gene ID:	3815		
UniProt:	P10721		
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin		
	Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, Production of		
	Molecular Mediator of Immune Response, Regulation of long-term Neuronal Synaptic Plasticity		
Application Details	Molecular Mediator of Immune Response, Regulation of long-term Neuronal Synaptic Plasticity		
Application Details Application Notes:	Molecular Mediator of Immune Response, Regulation of long-term Neuronal Synaptic Plasticity WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:200, ELISA(peptide) 1:20000-1:40000		
Application Notes:			
Application Notes: Restrictions:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:200, ELISA(peptide) 1:20000-1:40000		
	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:200, ELISA(peptide) 1:20000-1:40000		
Application Notes: Restrictions: Handling	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:200, ELISA(peptide) 1:20000-1:40000 For Research Use only		

Handling


	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. ABIN6267364 at 1/200 staining Rat kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunohistochemistry

Image 2. ABIN6267364 at 1/200 staining Human heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Western Blotting

Image 3. Western blot analysis of Phospho-KIT (Tyr703) expression in various lysates

Please check the product details page for more images. Overall 11 images are available for ABIN6255994.