antibodies.com

Datasheet for ABIN6256124 anti-YAP1 antibody (pSer127)

3 Images



Overview

Quantity:	100 µL	
Target:	YAP1	
Binding Specificity:	pSer127	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This YAP1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)	

Product Details

Immunogen:	A synthesized peptide derived from human YAP around the phosphorylation site of Serine 127
lsotype:	lgG
Specificity:	Phospho-YAP (Ser127) Antibody detects endogenous levels of YAP only when phosphorylated at Serine 127
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

Target:	YAP1
	Order at www.antibodies-online.com www.antikoerper-online.de www.anticorps-enligne.fr www.antibodies-online.cn
	International: +49 (0)241 95 163 153 USA & Canada: +1 877 302 8632 support@antibodies-online.com

Page 1/4 | Product datasheet for ABIN6256124 | 09/10/2023 | Copyright antibodies-online. All rights reserved.


Alternative Name:	YAP (YAP1 Products)
Background:	Description: Transcriptional regulator which can act both as a coactivator and a corepressor
	and is the critical downstream regulatory target in the Hippo signaling pathway that plays a
	pivotal role in organ size control and tumor suppression by restricting proliferation and
	promoting apoptosis (PubMed:17974916, PubMed:18280240, PubMed:18579750,
	PubMed:21364637). The core of this pathway is composed of a kinase cascade wherein
	STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and
	activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates
	and inactivates YAP1 oncoprotein and WWTR1/TAZ (PubMed:18158288). Plays a key role in
	tissue tension and 3D tissue shape by regulating cortical actomyosin network formation. Acts
	via ARHGAP18, a Rho GTPase activating protein that suppresses F-actin polymerization
	(PubMed:25778702). Plays a key role to control cell proliferation in response to cell contact.
	Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate
	cellular genes important for cell proliferation, cell death, and cell migration (PubMed:18158288)
	The presence of TEAD transcription factors are required for it to stimulate gene expression, cell
	growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction
	(PubMed:18579750).
	Gene: YAP1
Molecular Weight:	65kDa
Gene ID:	10413
UniProt:	P46937
Pathways:	MAPK Signaling, Stem Cell Maintenance, Regulation of Lipid Metabolism by PPARalpha
Application Details	
Application Notes:	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 $$ mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6256124 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. ABIN6267537 at 1/100 staining mouse kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunofluorescence (fixed cells)

Image 2. ABIN6267537 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Images

kDa	1	2
250-		
150-		
-00		
75-		
50-		-
37—		
25-		
20-		
15-		

Western Blotting

Image 3. Western blot analysis of YAP phosphorylation expression in COS7 whole cell lysates,The lane on the left is treated with the antigen-specific peptide.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN6256124 | 09/10/2023 | Copyright antibodies-online. All rights reserved.