antibodies -online.com







anti-DDIT3 antibody (pSer30)



Images



\sim	
()\/白	rview
OVC	

Quantity:	100 μL
Target:	DDIT3
Binding Specificity:	pSer30
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDIT3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human DDIT3 around the phosphorylation site of Ser30.
Isotype:	IgG
Specificity:	Phospho-DDIT3 (Ser30) Antibody detects endogenous levels of DDIT3 only when phosphorylated at Serine 30.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details


Target Details

Alternative Name:	DDIT3 (DDIT3 Products)
Background:	Description: Multifunctional transcription factor in ER stress response. Plays an essential role in
	the response to a wide variety of cell stresses and induces cell cycle arrest and apoptosis in
	response to ER stress. Plays a dual role both as an inhibitor of CCAAT/enhancer-binding protein
	(C/EBP) function and as an activator of other genes. Acts as a dominant-negative regulator of
	C/EBP-induced transcription: dimerizes with members of the C/EBP family, impairs their
	association with C/EBP binding sites in the promoter regions, and inhibits the expression of
	C/EBP regulated genes. Positively regulates the transcription of TRIB3, IL6, IL8, IL23,
	TNFRSF10B/DR5, PPP1R15A/GADD34, BBC3/PUMA, BCL2L11/BIM and ER01L. Negatively
	regulates, expression of BCL2 and MYOD1, ATF4-dependent transcriptional activation of
	asparagine synthetase (ASNS), CEBPA-dependent transcriptional activation of hepcidin (HAMP)
	and CEBPB-mediated expression of peroxisome proliferator-activated receptor gamma
	(PPARG). Inhibits the canonical Wnt signaling pathway by binding to TCF7L2/TCF4, impairing
	its DNA-binding properties and repressing its transcriptional activity. Plays a regulatory role in
	the inflammatory response through the induction of caspase-11 (CASP4/CASP11) which
	induces the activation of caspase-1 (CASP1) and both these caspases increase the activation
	of pro-IL1B to mature IL1B which is involved in the inflammatory response.
	Gene: DDIT3
Molecular Weight:	19kDa
Gene ID:	1649
UniProt:	P35638
Pathways:	Regulation of Muscle Cell Differentiation, ER-Nucleus Signaling, Skeletal Muscle Fiber
	Development, Cell RedoxHomeostasis
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %

Handling


	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images




Western Blotting

Image 1. Western blot analysis of CHOP phosphorylation expression in PMA treated Jurkat whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



Western Blotting

Image 2. Western blot analysis of Phospho-CHOP (Ser30) expression in various lysates



Immunohistochemistry

Image 3. ABIN6267488 at 1/100 staining human Breast carcinoma tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 5 images are available for ABIN6256253.