antibodies -online.com





Datasheet for ABIN6256330

anti-MAPT antibody (pSer422)






Go to Product page

\sim	
()\/\	view
	V I C V V

Overview	
Quantity:	100 μL
Target:	MAPT
Binding Specificity:	pSer422
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAPT antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
Immunogen:	A synthesized peptide derived from human Tau around the phosphorylation site of Ser422.
Isotype:	IgG
Specificity:	Phospho-Tau (Ser422) Antibody detects endogenous levels of Tau only when phosphorylated at Ser739, which site historically referenced as Ser422.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Target Details	
Target:	MAPT


Target Details

MAPT (MAPT Products)
Description: Promotes microtubule assembly and stability, and might be involved in the
establishment and maintenance of neuronal polarity. The C-terminus binds axonal
microtubules while the N-terminus binds neural plasma membrane components, suggesting
that tau functions as a linker protein between both. Axonal polarity is predetermined by
TAU/MAPT localization (in the neuronal cell) in the domain of the cell body defined by the
centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer
isoforms may preferentially play a role in its stabilization.
Gene: MAPT
80kDa
4137
P10636
MAPK Signaling, Microtubule Dynamics, M Phase, Regulation of Cell Size
WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
For Research Use only
Liquid
1 mg/mL
Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
glycerol.
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
should be handled by trained staff only.
-20 °C
Store at -20 °C. Stable for 12 months from date of receipt.
12 months




Western Blotting

Image 1. Western blot analysis of Tau phosphorylation expression in Mouse brain tissue lysates, The lane on the left is treated with the antigen-specific peptide.



Western Blotting

Image 2. Western blot analysis of Phospho-Tau (Ser422) expression in various lysates



Immunohistochemistry

Image 3. ABIN6267357 at 1/200 staining Human ganstric cancer tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 4 images are available for ABIN6256330.