antibodies -online.com





anti-WNK1 antibody (pThr60)





Go to Product page

Overview

Quantity:	100 μL
Target:	WNK1
Binding Specificity:	pThr60
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WNK1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details


Immunogen:	A synthesized peptide derived from human WNK1 around the phosphorylation site of Thr60.
Isotype:	IgG
Specificity:	Phospho-WNK1 (Thr60) Antibody detects endogenous levels of WNK1 only when phosphorylated at Threonine 60.
Predicted Reactivity:	Pig,Bovine,Sheep,Rabbit,Dog,Chicken
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

Target:	WNK1		
---------	------	--	--


Target Details

Alternative Name:	WNK1 (WNK1 Products)	
Background:	Description: Serine/threonine kinase which plays an important role in the regulation of electrolyte homeostasis, cell signaling, survival, and proliferation. Acts as an activator and	
	inhibitor of sodium-coupled chloride cotransporters and potassium-coupled chloride	
	cotransporters respectively. Activates SCNN1A, SCNN1B, SCNN1D and SGK1. Controls sodium	
	and chloride ion transport by inhibiting the activity of WNK4, by either phosphorylating the	
	kinase or via an interaction between WNK4 and the autoinhibitory domain of WNK1. WNK4	
	regulates the activity of the thiazide-sensitive Na-Cl cotransporter, SLC12A3, by	
	phosphorylation. WNK1 may also play a role in actin cytoskeletal reorganization.	
	Phosphorylates NEDD4L. Acts as a scaffold to inhibit SLC4A4, SLC26A6 as well as CFTR	
	activities and surface expression, recruits STK39 which mediates the inhibition (By similarity).	
	Gene: WNK1	
Molecular Weight:	230kDa	
Gene ID:	65125	
UniProt:	Q9H4A3	
Application Details		
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	




Western Blotting

Image 1. Western blot analysis of WNK1 phosphorylation expression in EGF treated 293 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6267692 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 3. Western blot analysis of Phospho-WNK1 (Thr58) expression in various lysates

Please check the product details page for more images. Overall 4 images are available for ABIN6256585.