antibodies -online.com





anti-CBL antibody (pTyr674)

3 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	CBL
Binding Specificity:	pTyr674
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CBL antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human CBL around the phosphorylation site of Tyr674.
Isotype:	IgG
Specificity:	Phospho-CBL (Tyr674) Antibody detects endogenous levels of CBL only when phosphorylated at Tyrosine 674.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Target Details

|--|


Target Details

Alternative Name:	CBL (CBL Products)
Background:	Description: Adapter protein that functions as a negative regulator of many signaling pathways
	that are triggered by activation of cell surface receptors. Acts as an E3 ubiquitin-protein ligase,
	which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to
	substrates promoting their degradation by the proteasome. Recognizes activated receptor
	tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, EGFR, CSF1R, EPHA8
	and KDR and terminates signaling. Recognizes membrane-bound HCK, SRC and other kinases
	of the SRC family and mediates their ubiquitination and degradation. Participates in signal
	transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast
	differentiation and apoptosis. Essential for osteoclastic bone resorption. The 'Tyr-731'
	phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to
	the cell membrane in a signaling pathway that is critical for osteoclast function. May be
	functionally coupled with the E2 ubiquitin-protein ligase UB2D3.
	Gene: CBL
Molecular Weight:	120kDa
Gene ID:	867
UniProt:	P22681
Pathways:	TCR Signaling, Interferon-gamma Pathway, EGFR Signaling Pathway, EGFR Downregulation, VEGFR1 Specific Signals
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling


	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images




Western Blotting

Image 1. Western blot analysis of CBL phosphorylation expression in Na2VO3 treated HepG2 whole cell lysates,The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6267436 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.



Immunohistochemistry

Image 3. ABIN6267436 at 1/100 staining Mouse intestine tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.