# antibodies -online.com





## anti-SMARCC2 antibody (Internal Region)

2 Images



Go to Product page

$\sim$	
( )\/△	rview
$\cup$	1 410 44

Target:

100 μL
SMARCC2
Internal Region
Human, Mouse, Rat
Rabbit
Polyclonal
This SMARCC2 antibody is un-conjugated
Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF),
Immunocytochemistry (ICC)
A synthesized peptide derived from human SMRC2, corresponding to a region within the
internal amino acids.
IgG
SMRC2 Antibody detects endogenous levels of total SMRC2.
Pig,Bovine,Horse,Sheep,Rabbit,Dog
The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling
Resin (Thermo Fisher Scientific).

SMARCC2


### **Target Details**


Alternative Name:	SMARCC2 (SMARCC2 Products)
Background:	Description: Involved in transcriptional activation and repression of select genes by chromatin
	remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin
	remodeling complexes that carry out key enzymatic activities, changing chromatin structure by
	altering DNA-histone contacts within a nucleosome in an ATP-dependent manner
	(PubMed:11018012). Can stimulate the ATPase activity of the catalytic subunit of these
	complexes (PubMed:10078207). May be required for CoREST dependent repression of
	neuronal specific gene promoters in non-neuronal cells (PubMed:12192000). Belongs to the
	neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-
	specific chromatin remodeling complex (nBAF complex). During neural development a switch
	from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons
	exit the cell cycle and become committed to their adult state. The transition from proliferating
	neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of
	the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into
	neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are
	exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C
	subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-
	renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along
	with CREST plays a role regulating the activity of genes essential for dendrite growth (By
	similarity). Critical regulator of myeloid differentiation, controlling granulocytopoiesis and the
	expression of genes involved in neutrophil granule formation (By similarity).
	Gene: SMARCC2
Molecular Weight:	160 kDa
Gene ID:	6601
UniProt:	Q8TAQ2
Application Details	
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL

#### Handling

Buffer:	Rabbit lgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

#### **Images**





#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6274439 staining HuvEc cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.

#### **Western Blotting**

**Image 2.** Western blot analysis of extracts from HUVEC cells using SMRC2 antibody.