# antibodies .- online.com





## anti-Meningioma 1 antibody (Internal Region)

2 Images

Target:




Go to Product page

Overview	
Quantity:	100 μL
Target:	Meningioma 1 (MN1)
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Meningioma 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human MN1, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	MN1 Antibody detects endogenous levels of total MN1.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	

Meningioma 1 (MN1)


## **Target Details**

Alternative Name:	MN1 (MN1 Products)
Background:	Description: Transcriptional activator which specifically regulates expression of TBX22 in the
	posterior region of the developing palate. Required during later stages of palate development
	for growth and medial fusion of the palatal shelves. Promotes maturation and normal function
	of calvarial osteoblasts, including expression of the osteoclastogenic cytokine
	TNFSF11/RANKL. Necessary for normal development of the membranous bones of the skull
	(By similarity). May play a role in tumor suppression (Probable).
	Gene: MN1
Molecular Weight:	135 kDa
Gene ID:	4330
UniProt:	Q10571
Application Details	
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### **Western Blotting**

**Image 1.** Western blot analysis of extracts from COLO cells, using MN1 antibody.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274324 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.