antibodies -online.com







anti-MYF5 antibody (N-Term)



Images



| () | 11/0 | r\ /1 | $\triangle 1 $ |
|-----|--------------------|-------|----------------|
| | $\lor \lor \vdash$ | 1 V I | ew |
| | | | |


| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | MYF5 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MYF5 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC) |
| Product Details | |
| | |

| Immunogen: | A synthesized peptide derived from human MYF5, corresponding to a region within N-terminal amino acids. |
|-----------------------|--|
| Isotype: | IgG |
| Specificity: | MYF5 Antibody detects endogenous levels of total MYF5. |
| Predicted Reactivity: | Bovine, Horse, Sheep, Rabbit |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific). |

Target Details

Target Details

| rarget Details | |
|---------------------|--|
| Alternative Name: | MYF5 (MYF5 Products) |
| Background: | Description: Acts as a transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation. Together with MYOG and MYOD1, co-occupies muscle-specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein. Gene: MYF5 |
| Molecular Weight: | 28 kDa |
| Gene ID: | 4617 |
| UniProt: | P13349 |
| Pathways: | Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development |
| Application Details | |
| Application Notes: | WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20 °C. Stable for 12 months from date of receipt. |
| Expiry Date: | 12 months |




Western Blotting

Image 1. Western blot analysis of extracts from HepG2 cells, using MYF5 antibody.



Immunofluorescence (fixed cells)

Image 2. ABIN6274168 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.



Western Blotting

Image 3. Western blot analysis of extracts of various tissue sample, using MYF5 Antibody .