antibodies -online.com







anti-RPS18 antibody

Images



\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Quantity:	100 μL
Target:	RPS18
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RPS18 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide
Isotype:	IgG
Specificity:	RPS18 Antibody detects endogenous levels of total RPS18
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	RPS18
Alternative Name:	RPS18 (RPS18 Products)
Background:	Description: Located at the top of the head of the 40S subunit, it contacts several helices of the

Target Details


	18S rRNA.
	Gene: RPS18
Molecular Weight:	17 kDa
Gene ID:	6222
UniProt:	P62269


Application Details

Application Notes:	WB 1:500~1:1000, IF/ICC 1:100-1:500
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months





Western Blotting

Image 1. Western blot analysis of extracts from K562 cells, using RPS18 antibody.



Immunofluorescence (fixed cells)

Image 2. ABIN6274750 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.