



[Go to Product page](#)

Datasheet for ABIN6257500
anti-ST5 antibody (Internal Region)

2 Images

Overview

Quantity:	100 µL
Target:	ST5
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ST5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human ST5, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ST5 Antibody detects endogenous levels of total ST5.
Predicted Reactivity:	Bovine,Horse,Sheep,Rabbit,Dog,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	ST5
---------	-----

Target Details


Alternative Name:	ST5 (ST5 Products)
Background:	<p>Description: Guanine nucleotide exchange factor (GEF) which may activate RAB9A and RAB9B. Promotes the exchange of GDP to GTP, converting inactive GDP-bound Rab proteins into their active GTP-bound form. May be involved in cytoskeletal organization and tumorigenicity.</p> <p>Isoform 1 seems to be involved in a signaling transduction pathway leading to activation of MAPK1/ERK2. Isoform 3 may block ERK2 activation stimulated by ABL1. Isoform 3 may alter cell morphology and cell growth.</p> <p>Gene: ST5</p>
Molecular Weight:	126 kDa
Gene ID:	6764
UniProt:	P78524

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of extracts from COLO205 cells, using ST5 antibody.



Immunofluorescence (fixed cells)

Image 2. ABIN6274303 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.